

## **New Records of Noctuidae (Lepidoptera) from the Korean Peninsula (II). Quadrifinae**

**AHN, Sung Bok<sup>1</sup>, Vladimir S. KONONENKO<sup>2</sup> and Kyu Tek PARK<sup>3</sup>**

<sup>1</sup>Division of Entomology, Agricultural Sciences Institute,  
Rural Development Administration, Suwon, 441-707, Korea

<sup>2</sup>Institute of Biology and Pedology, Far East Scientific Center,  
Russian Academy of Sciences, Vladivostok 22, RF60022, Russia

<sup>3</sup>Center for Insect Systematics, Kangwon National University, Chuncheon, 200-707, Korea

---

**Abstract** This list is based on the Noctuid materials of the collections of Agricultural Sciences Institute, Forest Research Institute, Center for Insect Systematics of Kangwon National University and Kyeongsang National University. A total of 24 species which belong to Noctuidae "Quadrifinae-group" are recorded from Korea for the first time. Among them, 12 species already known from Japan are reported for the first time from the continental Asia.

**Key words** Systematics, Fauna, Lepidoptera, Noctuidae, Quadrifinae, Korea

---

### **INTRODUCTION**

This paper, dealing with the Noctuid fauna of the Korean Peninsula is the second contribution of our study. This part contains a total of 24 species, belonging to the so called "Quadrifinae group" (Euteliinae, Chloephorinae, Plusiinae, Catocalinae, Ophiderinae, Hypenodinae, Rivulinae, Hypeninae) which are reported for the first time from Korea. Among them, twelve species are firstly reported for the continental Asia. Two subfamilies (Acontiinae and Herminiinae) are not included in this paper. For our study we used the materials preserved in the same specimen-rooms that previous paper dealt (Ahn *et al.*, 1994). The following abbreviations are used for the depositing places of collections: [ASI]-Agricultural Sciences Institute; [CIS]-Center for Insect Systematics; [FRI]-Forest Research Institute; [KSNU]-Kyeongsang National University. For the notes of distribution, we used abbreviation RFE which means Russian Far East region. Abbreviations of the provincial names for examined materials are given as GG-Gyeonggi Prov.; GW-Gangwon Prov.; CN-Chungnam Prov.; CB-Chungbuk Prov.; JN-Jeonnang Prov.; JB-Jeongbuk Prov.; GN-Gyeongnam Prov.; GB-Gyeongbuk Prov.; CJ-Cheju Prov. For the distribution range in Korea, four divisional words were used: Northern- northern part of the Korean Peninsula (upper area of DMZ, DPRK area); Central- central part of the Korean Peninsula (including GG, GW, CB and CN of ROK); Southern- southern part of the Korean Peninsula (including JB, JN, GB and GN of ROK); Cheju-Cheju province.

### **A LIST OF NEWLY ADDED SPECIES FROM THE KOREAN PENINSULA**

#### **Subfamily Euteliinae 비행기밤나방亞科**

*Eutelia grabczewskii* Püngeler 갈색점비행기밤나방 (Plate I-1)

*Eutelia grabczewskii* Püngeler, 1903, Deut. ent. Zeit. Iris 16: 289, pl. 6: 5.

**Material examined.** [ASI]-1 ♂, Mt. Naejangsan, JB, 11. VI. 1975 (J. -Y. Shim); 1 ♀, Mt. Pekunsan near Gwangyang, JN, 23. VI. 1991 (S. -B. Ahn); [CIS]-1 ♂, Gwangjeung, GG, 8. VII. 1992; 1 ♂, Chuncheon, GW, 21. VII. 1992 (K. -T. Park & B. -K. Byun).

**Distribution.** Korea (Central, Southern), Japan (Hokkaido, Honshu, Shikoku, Kyushu).

**Note.** Korean data is the first record for the continental Asia.

### Subfamily Sarrothripinae 겹질밤나방亞科

#### *Characoma ruficrura* (Hampson) 상수리겹질밤나방 (Plate I-2)

*Hypothripa ruficrura* Hampson, 1905, Ann. Mag. nat. Hist., (7) 16: 540.

**Material examined.** [ASI]-1 ♂, 1 ♀, Cheonan, CN, 2. IX. 1984 (D. -H. Kim).

**Distribution.** Korea (Central), Japan (Honshu, Shikoku, Kyushu), Indonesia (Borneo), N. India.

**Note.** This specimen was emerged from the galls of *Quercus* sp. made by a Cynipid wasp, *Trichagalma serratae* Ashmead (Cynipidae; Hymenoptera).

### Subfamily Chloephorinae 푸른밤나방亞科

#### *Earias erubescens* Staudinger 연홍색푸른밤나방

*Earias erubescens* Staudinger, 1887, In Romanoff, Mem. Lepid. 3: 175.

*Earias jezoensis* Sugi, 1982, In Inoue, H. et al., Moths of Japan, I: 797, II: 378, pl. 194: 41, 42, pl. 371: 1.

**Material examined.** [CIS]-12 specimens, Mt. Dorimsan, Chuncheon, GW; Mokuje (Mt. Seolaksan), GW; Gwangyang, JN; Sogeuimgang, GW.

**Distribution.** Korea (Central, Southern), Japan (Hokkaido, Honshu), RFE (from Amur basin to Primorye territory).

**Note.** The synonym cited was introduced from Sugi (1990), who compared the types of *E. erubescens* and *E. jezoensis* (Kononenko, 1990: 11, 30, No 219).

#### *Earias roseoviridis* Sugi 큰분홍무늬푸른밤나방 (Plate I-3)

*Earias roseoviridis* Sugi, 1982, In Inoue H. et al., Moths of Japan, I: 379, II: 797, pl. 194: 39, pl. 371: 3.

**Material examined.** [ASI]-1 ♀, Mt. Odaesan, GW, 23. V. 1980 (K. -T. Park); 1 ♂, Seomyon, Yangyang, GW, 24. V. 1987 (K. -T. Park); 1 ♂, Chuncheon, GW, 1. V. 1989 (K. -T. Park); 3 ♀, Seoguipo, CJ, 5. VII. 1986 (K. -T. Park).

**Distribution.** Korea (Central, Cheju), Japan (Amami-Oshima, Okinawa).

**Note.** Korean data is the first record of this species for the continental Asia.

#### *Paracrama angulata* Sugi 끝갈색푸른밤나방 (Plate I-4)

*Paracrama angulata* Sugi, 1985, Jap. Heteroc. J. 131: 90.

**Material examined.** [KSNU]-1 ♀, Geumgok (Jinyang), GN, 20. VI. 1982 (Y. -S. Liu).

**Distribution.** Korea (Southern), Japan (Honshu, Shikoku, Kyushu, Tsushima), Taiwan.

**Note.** This species was recorded as *P. dulcissima* Walker by Sugi (1982), but he corrected the scientific name as new one in 1985.

### Subfamily Plusiinae 은무늬나방亞科

#### *Abrostola abrostolina* (Butler, 1879) 활무늬알락밤나방 (Plate I-5)

*Inguridia abrostolina* Butler, 1879, Ann. Mag. nat. Hist. (5) 4: 354.

*Abrostola arensis* Leech, 1889 (nec Guenee, 1882), Proc. zool. Soc. Lond. 1889: 535.

*Abrostola tocionis* Marumo, 1917, Colln Essay Mr. Y. Nawa 3: 29, text fig.

**Material examined.** [ASI]-1 ♂, Gwangleung 26-30. VII. 1991 (K. -T. Park); [CIS]-1 ♂, Gwangleung, GG, 17. V. 1988 (K. -T. Park); 3 ♂, Gwangleung, GG, 8. VII 1989 (K. -T. Park & B. -K. Byun).

**Distribution.** Korea (Central), Japan (Honshu, Shikoku, Kyushu, Tsushima), China, Taiwan.

#### *Plusiopalpa adrasta* (Felder et Rogenhofer) 아시아은무늬밤나방 (Plate I-6)

*Plusiopalpa adrasta* Felder et Rogenhofer, 1874, Reise ost. Fregatte Novara (Zool.) 2 (Abt. 2): pl. 110, fig. 35.

**Material examined.** [ASI]-1 ♀, Mt. Seolaksan, GW, 9. IX. 1993 (S. -B. Ahn).

**Distribution.** Korea (Central), S. China, Vietnam, Indo-China, Indonesia (Borneo), India, SE Hymalaya.

**Note.** This species is distributed in the tropical regions of the old world. In Korea, this is probably a migrating species.

### Subfamily Catocalinae 뒷날개밤나방亞科

#### *Achaea janata* (Linnaeus) 흰별수중다리밤나방 (Plate I-7)

*Phalaena janata* Linnaeus, 1758, Syst. Nat. (Ed. 10) 1: 527.

*Noctua melicerta* Drury, 1773, Illust. nat Hist. 1 (1770): 46, pl. 23: 1.

*Noctua tigrina* Fabricius, 1781, Spec. Ins. 2: 218.

*Catocala traversii* Fereday, 1877, Trans. N. Z. Inst. 9: 457.

*Ophiura ekeikei* Beth. -Baker, 1906, Nov. Zool. 13: 256.

**Material examined.** [ASI]-1 ♀, Suwon, GG, 20. IX. 1958; 1 ♀, Cheju, CJ, 20. VI. 1975 (H. -S. Kim); 1 ♀, Seongpanak, CJ, 2, VIII. 1984 (K. -S. Lee).

**Distribution.** Korea (Central, Cheju), Japan (from Honshu to Okinawa), China, Taiwan, Indo-China, India, Philippines, New Guinea, Micronesia, Oceania, Australia.

**Note.** The species is widely distributed in the Indo-Australian region and SE Asia.

### Subfamily Ophiderinae 짙름나방亞科

#### *Lygephila cracca* (Denis et Schiffermüller) 잿빛목검은밤나방 (Plate I-8)

*Noctua cracca* Denis et Schiffermüller, 1775, Ankun. Syst. Werk. Schmett. Wien. 1775: 94.

**Material examined.** [ASI]-1 ♂, Hongcheon, GW, 28. VI. 1989 (K. -S. Lee).

**Distribution.** Korea (Central), Japan (Hokkaido), RFE (from Amur region to Primorye and SaGhalin), Siberia, Central Asia, Europe.

**Note.** This is a widely distributed transpalearctic species.

#### *Lygephila subrecta* Sugi 극동목검은밤나방 (Plate II-1)

*Lygephila subrecta* Sugi, 1982, In Inoue H. et al., Moths of Japan I: 857, II: 393, Pl. 210: 20, 21; 376: 1, 3.

**Material examined.** [ASI]-1♂, Hongcheon, GW, 20. V. 1988 (P. -K. Choi); 1♂, Suwon, GG, 23. VII. 1980 (C. -G. Yoo); 2♂, Suwon, 23, 26. VIII. 1976 (K. -B. Uhm); 1♀, Suwon, 22. VI. 1976 (K. -T. Park); 1♀, Suwon, 6. V. 1989 (S. -B. Ahn); [CIS]-3♂, 3♀, Suwon, 6. VI. 1989; 22. VI. 1976; 23, VII. 1980; 27. VI. 1976 (B. -K. Byun, K. -T. Park, C. -G. Yoo).

**Distribution.** Korea (Central), Japan (Hokkaido, Honshu).

**Note.** Korean data is the first record for the continental Asia.

**Anomis longipennis Sugi 붉은잎밤나방 (Plate II-2)**

*Anomis longipennis* Sugi, 1982, In Inoue H. *et al.*, Moths of Japan I: 853, II: 393, pl. 211: 12-14, pl. 367: 10, 15, 19.

**Material examined.** [FRI]-5♂, 3♀, Gwangleung, GG, 6. XI. 1990 (G. -J. Weon); [KSNU]-1♂, Namhae (Idongmyon., Sinjonri, Mt. Geumsan), GN, 28-29. VII. 1990 (D. -S. Ku).

**Distribution.** Korea (Central, Southern), Japan (Hokkaido, Honshu, Shikoku., Kyushu).

**Note.** Korean data is the first record for the continental Asia.

**Daddala lucilla (Butler) 톱날무늬잎밤나방 (Plate II-3)**

*Sypna lucilla* Butler, 1881, Trans. Ent. Soc. Lond. 1881: 206.

**Material examined.** [ASI]-1♀, Namhae (Mt. Geumsan), GN, 24. VII. 1985 (K. -T. Park); 1♀, Seongpanak, CJ, 2. VIII. 1984 (K. -S. Lee).

**Distribution.** Korea (Southern, Cheju), S. Japan (from Honshu to Okinawa and sothern island), China, RFE (S. Primorye, as a migrant), Taiwan, Burma, Borneo, Sumatra, Java, India, New Guinea.

**Wilemaniella angulata (Wileman) 우묵날개팔름나방 (Plate II-4)**

*Rivula angulata* Wileman, 1911, Trans. ent. Soc. Lond. 1911: 267.

**Material examined.** [ASI]-2♂, Mt. Yumyongsan, GG, 30. VII. 1990 (S. -H. Oh); 1♂, 1ex., Mt. Pekunsan near Gwangyang, JN, 25. VI. 1991 (S. -B. Ahn).

**Distribution.** Korea (Central, Southern), Japan (Honshu, Shikoku, Kyushu, Tsushima).

**Note.** Korean data is the first record for the continental Asia.

**Paragabara secunda Remm 극동팔름나방 (Plate II-5)**

*Paragabara secunda* Remm, 1983, Entomol. Obozr. 62: 596: 7.

**Material examined.** [CIS]-1♀, Bongmyongri, Hongcheon, GW, 30. VI. 1992 (K. -T. Park).

**Distribution.** Korea (Central), RFE (S. Primorye).

**Microxyla confusa (Wileman) 세모팔름나방 (Plate II-6)**

*Rivula confusa* Wileman, 1911, Trans. ent. Soc. Lond. 1911: 268.

**Material examined.** [ASI]-2♀, Mt. Yumyongsan, GG, 30. VII. 1990 (S. -H. Oh); 1ex., Suwon, GG, 15. VII. 1976 (K. -T. Park); 1♂, 1♀, Mt. Pekunsan near Gwangyang, JN, 25. VI. 1991 (S. -B. Ahn); 1♂, 1♀, Cheju, CJ, 5. VII. 1986 (K. -T. Park).

**Distribution.** Korea (Central, Southern, Cheju), Japan (Honshu, Shikoku, Kyushu, Tsushima, Yakushima).

**Note.** Korean data is the first record for the continental Asia.

**Prolophota trigonifera Hampson 두점깨다시팔름나방 (Plate II-7)**

*Prolophota trigonifera* Hampson, 1896, Fauna Br. India (Moths) 4: 547, fig 284.

**Material examined.** [ASI]-1♀, Chuncheon, GW, 24. VII. 1991 (S. -B. Ahn); [CIS]-1♂, Mt. Seola-

ksan, GW, 10. VIII. 1992 (K. -T. Park & B. -K. Byun).

*Distribution.* Korea (Central), Japan (Honshu, Shikoku, Kyushu, southern islands), Sri Lanka.

***Gonepatica butleri* (Leech) 볼록팔름나방 (Plate II-8)**

*Camptochilus butleri* Leech, 1900, Trans. ent. Soc. Lond. 1900: 647.

*Material examined.* [ASI]-1 ♂, Yangyang, GW, mid. V. 1987 (K. -T. Park); 1 ♀, Muju, JB, 12. VIII. 1975 (K. -T. Park); [CIS]-1 ex., Chuncheon, GW, 22. V. 1985 (K. -T. Park); 1 ex., Chuncheon, GW, 21. VII. 1992 (K. -T. Park & B. K. Byun); 1 ex., Chuncheon dam, GW, 12. VIII. 1991 (K. -T. Park); 1 ex., Mt. Gyeongbongsan, GW, 2. VIII. 1989 (K. -T. Park); 2 exs., Yangyang, GW, 10. V. 1987 (K. -T. Park); 1 ex., Sogyeunggang, GW, 23. V. 1988 (K. -T. Park).

*Distribution.* Korea (Central, Southern), Japan (Honshu, Shikoku, Kyushu, Tsushima, Yakushima).

*Note.* Korean data is the first record from the continental Asia.

***Anachrostis nigripunctalis* (Wileman) 점무늬팔름나방 (Plate III-1)**

*Rivula nigripunctalis* Wileman, 1911, Trans. ent. Soc. Lond. 1911: 268.

*Material examined.* [ASI]-5 ♂, 3 ♀, Suwon, GG, 2, 23, 28. VIII. 1974 (K. -T. Park); 2 ♂, Suwon, GG, 7. VI., 16. IX. 1976 (K. -T. Park); 1 ♂, Yangyang, GW, 26. VII. 1980 (K. -R. Choi); 1 ♂, Pyeongchang, GW, 24. VI. 1988 (S. -H. Oh); 1 ♂, Hongcheon, GW, 6. VII. 1987 (K. -S. Lee); 1 ♀, Suanbo, CB, 1. VII. 1990 (S. -H. Oh); 2 ♂, Mt. Gyeryongsan, CN, 20. VI. 1980 (K. -R. Choi); 2 ♂, Onyang, CN, 26. VI. 1990 (S. -H. Oh); 1 ♀, Mt. Jirisan, JB, 17. VII. 1976 (Y. -Y. Ha); 1 ♂, 1 ♀, Mt. Naejangsan, JB, 16. VI. 1992 (S. -B. Ahn); 15 ♂, 4 ♀, Mt. Pekunsan (Gwangyang), JN, 25. VI. 1991 (S. -B. Ahn); 1 ♂, Mt. Mudeungsan, JN, 29. VI. 1990 (S. -H. Oh); [CIS]-27 exs., Bongmyongri, Hongcheon, GW; Mt. Palbongsan, GW; Chuncheondam, GW; Honcheon, GW; Soyang dam, GW; Chuncheon, GW; Topyong, CJ; Gwaneumsa, CJ, 9. VI. -31. VIII. 1986-1992 (K. -T. Park and B. -K. Byun, S. -G. Lee).

*Distribution.* Korea (Central, Southern), Japan (from Hokkaido to Kyushu).

*Note.* Korean data is the first record for the continental Asia.

***Anachrostis minutissima* Sugi 작은점무늬팔름나방 (Plate III-2)**

*Anachrostis minutissima* Sugi, 1982, In Inoue H. *et al.*, Moths of Japan. 1: 890, 2: 402, pls. 219: 40, 41, 38: 3, 4.

*Material examined.* [CIS]-11 exs., Mt. Soyosan, GG; Mt. Seolaksan, GW; Mt. Odaesan, GW; Mt. Gyeongbongsan, GW; Mt. Samaksan, GW; Chuncheon, GW; Gotan, GW; Hongcheon, GW, 6. VIII. -25. IX; 3. XI, 1987-1992 (K. -T. Park, B. -K. Byun).

*Distribution.* Korea (Central), Japan (Yakushima, Nishi-Oshima Is.).

*Note.* This smallest moth has been known only from the small islands of southern part of Japan. Korean data is the first report for the continental Asia

**Subfamily Hypenodinae 꼬마팔름나방亞科**

***Hypenodes rectifascia* Sugi 흰줄꼬마팔름나방 (Plate III-3)**

*Hypenodes rectifascia* Sugi, 1982, In Inoue H. *et al.*, Moths of Japan I: 899, II: 403, pl. 220: 21, pl. 379: 11, pl. 380: 12.

*Material examined.* [ASI]-1 ♀, Suwon, GG, 20. V. 1976 (K. -T. Park); 1 ♀, Suwon, GG, 23. VII. 1975 (K. -T. Park); 2 ♂, Suwon, GG, 10. VIII. 1975, 30. VIII. 1976 (K. -T. Park); 5 ♂, 3 ♀, Suwon, GG, 13. IX. 1975 (K. -T. Park); 1 ♀, Gwangleung, GG, 10. VII. 1982 (K. -T. Park); 1 ♀,

Mt. Yumyongsan, GG, 30. VII. 1990 (S. -H. Oh); [CIS]-1 ♂, Mt. Soyosan, GG, 3. XI. 1992 (K. -T. Park); 1 ♂, Gwangleung, GG, 27. VII. 1988 (K. -T. Park); 1 ♂, Chuncheon, GW, 10. VII. 1989 (K. -T. Park).

*Distribution.* Korea (Central), Japan (Honshu, Kyushu).

*Note.* Korean data is the first record for the continental Asia.

***Schrankia kogii* Inoue 동근점꼬마팔름나방 (Plate III-4)**

*Schrankia kogii* Inoue, 1979, Tinea 10: 304, fig. 5, 11.

*Material examined.* [ASI]-1 ♀, Yongin, GG, 28. V. 1989 (K. -S. Lee); 1 ♀, Suwon, GG, 16. IX. 1976 (K. -T. Park); 1 ♂, 1 ♀, Suwon, GG, 10. IX. 1974 (Y. -I. Lee); 2 ♀, Gwangleung, GG, 10. VII. 1982 (K. -T. Park); [CIS]-1 ♀, Chuncheon, GW, 9. IX. 1988 (K. -T. Park) and further specimens from Mt. Yumyongsan, GG.

*Distribution.* Korea (Central), Japan (Hokkaido, Honshu).

*Note.* Korean data is the first record for the continental Asia.

***Schrankia masuii* Inoue 연갈색꼬마팔름나방 (Plate III-5)**

*Schrankia masuii* Inoue, 1979, Tinea 10: 303, figs 1, 9.

*Material examined.* [ASI]-1 ♂, Suwon, GG, 19. VI. 1985 (J. -O. Lee); 4 ♂, 1 ♀, Suwon, GG, 18. V. 1977, 11. VI. 1974, 11. VI., 18. IX. 1974 1976 (K. -T. Park); 1 ♀, Suwon, GG, 18. VII. 1982 (D. -J. Im); 1 ♂, Mt. Weolaksan, CB, 20. VI. 1984 (S. -B. Ahn).

*Distribution.* Korea (Central), Japan (Honshu, Shikoku, Kyushu).

*Note.* The Korean data is the first report for the continental Asia.

***Schrankia costaestrigalis* (Stephens) 검은줄꼬마팔름나방 (Plate III-6)**

*Cledeobia costaestrigalis* Stephens, 1843, Ill. Br. Ent. (Haustellata) 4: 21.

*Material examined.* [CIS]-1 ♂, Mt. Samaksan, 21. XI. 1989 (K. -T. Park); [ASI]-9 exs., Mt. Myongjisan, GG; Isl. Seonban, GG; Mt. Gyeongbansan, GW; Mt. Soyosan, GG; Chuncheon, GW; Gwangyang, JN; Cheju N. Univ., 17-18. V; 4. IV; 8-28. VII; 24. VIII; 3. IX; 21. IX. 10, 1989-92 (K. -T. Park, B. -K. Byun, S. -H. Oh & H. -Y. Choi).

*Distribution.* Korea (Central, Southern, Cheju), Japan (from Honshu to Okinawa), RFE (Primorye), SE Asia region, Middle East, Central and South Europe, Australia.

**Subfamily Rivulinae 노랑팔름나방亞科**

***Rivula unctalis* Staudinger 끝갈색노랑팔름나방**

*Rivula unctalis* Staudinger, 1892, In Romanoff, Mem. Lepid. 6: 631.

*Material examined.* [CIS]-26 exs., Mt. Myongjisan, GG; Chuncheon, GW; Bongmyongri, Hongcheon, GW; Mt. Myongjisan, GW; Gwangleung, GG; Mt. Daedunsan, CN; Cheju Nat. Univ., CJ; 16. V.-3. VI; 8. VII.-2. IX. 1988-1992 (K. -T. Park, B. -K. Byun, G. -J. Weon).

*Distribution.* Korea (Central, Cheju), Japan (Hokkaido), RFE (from Amur Basin to Primorye).

**ACKNOWLEDGEMENTS**

We appreciate to Agricultural Sciences Institute, Forest Research Institute, Kangwon National University and Kyeongsang National University for their kind allowance to the authors to use various facilities and financial supports of the Rural Development Administration. We would like to express

our thanks to Dr. Kui-Moon Choi and Mr. Man-Jong Han, Division of Entomology, Agricultural Sciences Institute for their kind helps during this study. Our thanks are also expanded to Professor Jung-Suk Park, Kyeongsang National University for admitting the examination of his collections. Our special thanks are expanded to Drs. L. Ronkay and L. Peregovits, Hungarian Natural History Museum and Professor Yoo-Hang Shin, Kyunghee University for the fruitful discussion and critical revision of the manuscript.

## REFERENCES

- Ahn S. B., L. Ronkay and K. T. Park. 1994. Faunistic data of Noctuidae (Lepidoptera) from Cheju-do, Southern Island of the Korean Peninsula. *Insecta Koreana* (suppl.): (in press).
- Ahn S. B., V. S. Kononenko and K. T. Park. 1994. New Records of Noctuidae (Lepidoptera) from the Korean Peninsula. (I) Trifinae. *Insecta Koreana* 11: (in press).
- Kononenko, V. S. 1990. Synonymic Check List of the Noctuidae of the Primorye Territory, the Far East of U.S.S.R. *Tinea* 13, Suppl. 40 pp.
- Sugi, S. 1982. Noctuidae. In Inoue, H. *et al.*, *Moths of Japan*, 1: 669-936, 2: 344-409, pls. 163-226, 355-392.
- Sugi, S. 1985. Three new species of Noctuidae: corrections to misidentifications found in *Moths of Japan Jap. Heterocerists' J.* 131: 89-91.
- Sugi, S. 1990. *Earias erubescens* Staudinger, the valid name replacing *E. jejoensis* Sugi (Noctuidae, Chloephorinae). *Japan Heterocerists' J.* 159: 154.
- Sviridov, A. V. 1990. Five new for the USSR species of noctuid moths from Primorye. *Vestn. zoologii*. No. 2: 87

## 한국産 밤나방科(나비目)의 未記錄種 (II). Quadrifinae

安聖復<sup>1</sup> · V. S. Kononenko<sup>2</sup> · 朴奎澤<sup>3</sup>

<sup>1</sup>農村振興廳 農業技術研究所 昆蟲科

<sup>2</sup>러시아科學院, 極東土壤生物研究所, 블라디보스톡, 러시아

<sup>3</sup>江原大學校 昆蟲系統分類研究센터


農業技術研究所, 林業研究院, 江原大學校 昆蟲系統分類研究센터, 慶尚大學校 生物科學에 所藏된 밤나방科 標本을 同定한 結果 확인된, "Quadrifinae" (Euteliinae, Sarothripinae, Chloephorinae, Plusiinae, Catocalinae, Ophiderinae, Hyphenodinae, Rivulinae and Hypheninae)에 속하는 韓國 未記錄種 24種을 報告한다. 이중 12種은 일본에서만 기록되어 있으나, 아시아 내륙에서는 처음 기록된다.

검색어: 분류, 나비목, 밤나방, 미기록종, Quadrifinae, 분포, 한국

(Received: March 30, 1994)

(Accepted: May 20, 1994)






**PLATE I**

1. *Eutelia grabczewskii* Püngeler
2. *Characoma ruficirra* (Hampson)
3. *Earias roseoviridis* Sugi
4. *Paracrama angulata* Sugi

5. *Abrostola abrostolina* (Butler)
6. *Plusiopalpa adrasta* (Felder et Rogenhofer)
7. *Achea janata* (Linnaeus)
8. *Lygephila cracca* (Denis et Schiffermüller)






# **PLATE II**

1. *Lygephila subrecta* Sugi
2. *Anomis longipennis* Sugi
3. *Daddala lucilla* (Butler)
4. *Wilemaniella angulata* (Wileman)

5. *Paragabara secunda* Remm
6. *Mycroxyla confusa* (Wileman)
7. *Prolophota trigonifera* Hampson
8. *Gonepatica butleri* (Leech)



**PLATE III**

1. *Anachrostitis nigripunctalis* (Wileman)
2. *Anachrostitis minutissima* Sugi
3. *Hypenodes rectifascia* Sugi

4. *Schrankia kogii* Inoue
5. *Schrankia masuii* Inoue
6. *Schrankia costaestrigalis* (Stephens)